Hydrology

Presented by

Chris Milly

① Land model LM3, used in all GFDL CMIP5 streams, contains new physical/hydrologic features that

- improve fluxes to atmosphere and oceans;

- support terrestrial biogeochemical modeling;

- represent impact-relevant hydrologic variables (streamflow, lake levels, water table).

② A focus on potential evapotranspiration (PET) helps reconcile conflicting projections of water availability and drought.

soil-veg
column
(tile)

landscape unit

lake tile

grid cell

river network

LM3 storage variables compare well with observations.

annual range of water storage (mm)

LM3 Hydrology: Evaluation

LM3 reproduces permafrost extent well.

permafrost extent

DICE: Diurnal Land/Atmosphere Coupling Experiment

(credit: Adrian Lock, Martin Best)

Work in Progress: Toward LM4

Hill-slope tiling (Subin)   Urban tiling

+ Higher Resolution
+ Biogeochemistry (Shevliakova)

Potential Evapotranspiration

Many hydrologic impact analyses have PET as a pivotal link:

Potential Evapotranspiration

- In a case study, PET change had been overestimated by a factor of ~3 vs. that implicit in the climate model.

- Substantial negative bias in runoff change results.

- SREX→AR5: "AR4 conclusions regarding global increasing drought since the 1970s were probably overstated."

Relative Change in PET

y = 3.0x

hydrologic model

climate model

(after Milly and Dunne [2011])

① Land model LM3, used in all GFDL CMIP5 streams, contains new physical/hydrologic features that

- improve fluxes to atmosphere and oceans;

- support terrestrial biogeochemical modeling;

- represent impact-relevant hydrologic variables (streamflow, lake levels, water table).

② A focus on potential evapotranspiration (PET) helps reconcile conflicting projections of water availability and drought.